

FIG. 1




FIG. 2A




FIG. 2B





FIG. 2C






FIG. 4A




FIG. 4B




FIG. 4C




FIG. 5




FIG. 6




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 12A




FIG. 12B




FIG. 12C




FIG. 13A




FIG. 13B




FIG. 13C




FIG. 14A




FIG. 14B




FIG. 15A




FIG. 15B




FIG. 15C

SERUM IgG ANTI-POLIOVIRUS IN
IMMUNIZED PIGTAIL MACAQUE




FIG. 16

pT7-IC:




FIGURE 17A

pT7-IC-Pr55^{gag}:




FIGURE 17B

pT7-IC-Pr55^{gag} (VP4/2A):




FIGURE 17C




FIGURE 18A

FIGURE 18B

09564 0199
4050 499960




FIGURE 19A

Phosphorimagery Quantitation of Samples
Analyzed by Northern Blot

Samples	Values
pT7-IC-Pr55 ^{gag}	19,062
pT7-IC-Pr55 ^{gag} (VP4/2A)	18,430
pT7-IC-Gag 1	98,800

FIGURE 19B




FIGURE 20



[illegible]

FIGURE 21A



pT7-IC




FIGURE 23A




FIGURE 23B

pT7-IC-CEA sig-




FIGURE 23C




FIGURE 25A

Anti-3Dpol

FIGURE 25B

Anti-CEA

FIGURE 25C

[³H] Leucine




FIGURE 26A




FIGURE 26B